

**EMMA ID: 05506**

**Gene: *Vmp1 / tmem49***

**Common name: VMP1 KO cond**

## Genotyping Information

Genotyping by end-point PCR based on gel is composed of a genespecific short range PCR using primers on wild type allele and a mutant allele-specific short range PCR. The combined results show the genotype of the mice. For example: mutant positive, wild type positive = Heterozygous.

### PCR primer pairs and expected size bands

| Assay        | Forward Primer | Reverse Primer | Expected Size Band (bp) |
|--------------|----------------|----------------|-------------------------|
| control-band | Primer 1459_27 | Primer 1459_28 | 266                     |
| Mutant       | Primer 1459_27 | Primer 1459_28 | 391                     |

### Primer sequences

| Primer Name    | Sequence 5' --> 3'   |
|----------------|----------------------|
| Primer 1459_27 | GCTTGCTGTGAATGGTTACC |
| Primer 1459_28 | TCAGATCAGCCTTCTGTAGG |

### PCR setup (Qiagen, Hot Start Plus)

| Component                     | Volume ( $\mu$ l) 1x | Final conc. |
|-------------------------------|----------------------|-------------|
| DNA (~ 50-100 ng)             | 2                    |             |
| Q-Solution (5x)               | 2,5                  | 0,5         |
| PCR-Buffer (10x)              | 2,5                  | 1           |
| DNTP mix (10 mM)              | 0,5                  | 0,2         |
| MgCl <sub>2</sub> (25 mM)     | 1,5                  | 1,5         |
| Primer 1 (10 pmol/ $\mu$ l)   | 1                    | 0,4         |
| Primer 2 (10 pmol/ $\mu$ l)   | 1                    | 0,4         |
| Taq Polymerase (5 U/ $\mu$ l) | 0,3                  | 0,06        |
| H <sub>2</sub> O*             | 13,7                 |             |
| Final volume                  | 25                   |             |


\* The amount of H<sub>2</sub>O is adjusted with the number of primer.

### Amplification conditions

| PCR Settings                                       | Temperature (°C)     | Time                       | # of cycles |
|--|----------------------|----------------------------|-------------|
| 1 Denaturation<br>(Melting)                        | 95°C                 | 5 min                      | 1           |
| 2 Amplification<br>(Melting, Annealing,<br>Polym.) | 94°C<br>60°C<br>72°C | 30 sec<br>45 sec<br>45 sec | 39          |
| 3 Polymerisation                                   | 72°C                 | 10 min                     | 1           |
| 4 Cooling  | 4°C                  | hold                       | 1           |

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.

### Gel Image



Separated by gel electrophoresis on a 2% agarose gel.