



Genotyping protocol

Afm

IR00003323 / E229

(ICS internal reference)

This report has been prepared by:

Valérie Rousseau
genotyping@igbmc.fr

This report has been validated by:

Sylvie Jacquot, PhD, Head of Genotyping Service
33 (0)3 88 65 57 44
genotyping @igbmc.fr

The first version of this report was finalized the: 07 Dec 2012

For any question, please contact:

Institut Clinique de la Souris - ICS - Mouse Clinical Institute
1 rue Laurent Fries, BP 10142
67404 Illkirch Cedex, France
Email: genotyping@igbmc.fr
Web site: <http://www-mci.u-strasbg.fr/>

TABLE OF CONTENTS


Table of contents	2
1. Genotyping protocol and data	2
1.1. Genotyping strategy.....	2
1.2. PCR protocol.....	4
1.3. Picture of genotyping with various alleles.....	4
2. Cre and Flp genotyping method.....	5
2.1. Cre and Flp genotyping.....	5
2.2. PCR Protocol.....	6

1. Genotyping protocol and data

This section describes the condition used at the Mouse Clinical Institute (ICS) to genotype your Afm Constitutive Knockout / Conditional Knockout (KO-cKO) project.

1.1. Genotyping strategy

The map below describes the position of the primers used for genotyping for each possible allele.



Sequence of primers used for genotyping:

Position	Primers	Sequence
Ef	5661	ACCCAGCCAAAGAGAACAAACAAAGG
Ef ²	5662	TCTCGTAGAAGTGAATGGACCACCC
Er	5666	GCTCGTTGGAAATGGATTCTGGGA
Kr	3277	CTCCTACATAGTGGCAGTGTGTTGGG
L3f	5663	GTGGAGCCCTTATAAAGTTGGACCC
L3r	5664	AGAGGGGTCACGATCCACAGGTT
L3r ²	5665	GGTTACCCCTCGGTGACGCATT
Lxr	3255	ACTGATGGCGAGCTCAGACCATAAC

² : For a selected position, a second primer was designed

PCR fragments expected size (bp):

Region analyzed	Primers used	Position on the primer <i>(see the map above)</i>	Targeted allele (KO allele) (L3)	cKO allele (L2)	KO allele (L-)	WildType allele (WT)
5' part of the selection marker	5661-3277	Ef / Kr	425	---	---	---
Presence of the distal loxP	5663-5664	L3f / L3r	396	396	---	443
Distal loxP specific PCR	5663-3255	L3f / Lxr	282	282	---	---
Excision of the selection marker	5661-5666	Ef / Er	7461*	557	---	416
Excision of the floxed exon(s), i.e. knock out	5662-5665	Ef ² / L3r ²	10000*	3118*	606**	3024**

*: this PCR product will not be observed using our PCR genotyping conditions (see description below)

**: this PCR is only verified if mice are generated

---: no Amplicon should be obtained

1.2. PCR protocol

This section describes the composition of the mix and cycling conditions used for genotyping.

Reagents:

- FastStart PCR Master (Roche)
- DNA (50ng/ μ l)
- 5' primer (100 μ M)
- 3' primer (100 μ M)
- Sterile H₂O

Volume:

- 7.5 μ l
- 1.5 μ l
- 0.06 μ l
- 0.06 μ l
- up to 15 μ l

Cycling conditions:

Temp	Time	#Cycles
95°C	4min	1
94°C	30s	
62°C	30s	34
72°C	1min	
72°C	7min	1
20°C	5min	1

NB: These PCR conditions have been optimized for high-throughput genotyping. Adaptation to small-scale may be required.


2. Cre and Flp genotyping method

The protocol used to segregate the cre and/or flp transgene is indicated below.

Detection of cre transgene and flp transgene is done using a multiplex assay: primer pairs were designed for each gene and for a positive control (Cpxm1 gene).

2.1. Cre and Flp genotyping

Schematic representation of the genotyping strategy



Sequence of primers used for genotyping:

Primers	Sequence
4029	ACTGGGATCTTCGAACCTTTGGAC
4030	GATGTTGGGGCACTGCTCATTACC
4045	CCATCTGCCACCCAGCCAG
4046	TCGCCATCTCCAGCAGG
4195	TCTTAGCGCAAGGGGTAGGATCG
4196	GTCCTGGCACGGCAGAAC

PCR fragments expected size (bp):

Primer pair	4045-4046	4195-4196	4029-4030
Region analyzed	Middle part of Cre transgene	Middle part of Flp transgene	Cpxm1 control gene
Control gene	/	/	397
Tg allele	281	328	/

2.2. PCR Protocol

This section describes the composition of the mix and cycling conditions used for genotyping.

Reagents	Volume
FastStart PCR Master (Roche)	7.5µl
DNA (50ng/µl)	1.5µl
5' primer (100 µM)	0.05µl
3' primer (100 µM)	0.05µl
Sterile H ₂ O	up to 15 µl

Cycling conditions are identical to those described in chapter 1.2