

**EMMA ID: 10141**

**Gene:** *Uqcrh*

**Common name:** *EPD0378\_3\_C07*

**Allele:** *Uqcrh* <sup>tm1b(EUCOMM)Wtsi</sup>

## Allele Information

Further information about the allele can be found on IMPC website at (copy the link to web browser)  
[https://www.mousephenotype.org/data/alleles/MGI:1913826/tm1b\(EUCOMM\)Wtsi](https://www.mousephenotype.org/data/alleles/MGI:1913826/tm1b(EUCOMM)Wtsi)

## Links to the general information

About IKMC resource

<https://www.infrafrontier.eu/knowledgebase/protocols/ikmc-products>

IKMC allele types

<http://www.i-dcc.org/kb/entry/89/>

Allele conversion guide - genotyping tm1b, tm1c and tm1d mice (assays infos available when required)

<http://www.mousephenotype.org/about-ikmc/targeting-strategies>

IMPC mouse phenotype data, search by the gene name

<http://www.mousephenotype.org/>

## Genotyping Information

Genotyping by end-point PCR based on gel is composed of a gene-specific short range PCR using primers on wild type allele and a mutant allele-specific short range PCR. The combined results show the genotype of the mice.

For example: mutant positive, wild type positive = Heterozygous.

### PCR primer pairs and expected size bands

Assay	Forward Primer	Reverse Primer	Expected Size Band (bp)
Mutant	Uqcrh 5'arm	LAR3	518
Wildtype	Uqcrh 5'arm	Uqcrh 3'arm	652

### Primer sequences

Primer Name	Sequence 5' --> 3'
Uqcrh 5'arm	aacttggactcctggtcctct
Uqcrh 3'arm	cttgcctcccccagaactagct
LAR3	CAACGGGTTCTTGTAGTCC

### PCR setup (Qiagen, Hot Start Plus)

Component	Volume ( $\mu$ l) 1x	Final conc.
DNA (~ 50-100 ng)	2	
Q-Solution (5x)	2,5	0,5
PCR-Buffer (10x)	2,5	1
DNTP mix (10 mM)	0,5	0,2
MgCl <sub>2</sub> (25 mM)	1,5	1,5
Primer 1 (10 pmol/ $\mu$ l)	1	0,4
Primer 2 (10 pmol/ $\mu$ l)	1	0,4
Taq Polymerase (5 U/ $\mu$ l)	0,3	0,06
H <sub>2</sub> O*	13,7	
Final volume	25	


\* The amount of H<sub>2</sub>O is adjusted with the number of primer.

### Amplification conditions

PCR Settings	Temperature (°C)	Time	# of cycles
1 Denaturation (Melting)	95°C	5 min	1
2 Amplification (Melting, Annealing, Polym.)	94°C 65°C 72°C	30 sec 45 sec 45 sec	39
3 Polymerisation	72°C	10 min	1
4 Cooling	12°C	hold	1

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.

### Gel Image



Separated by gel electrophoresis on a 2% agarose gel.

## Genotyping using PCR-assays for cassette detection

LacZ reporter, Neo selection cassettes are inserted into the Knockout-first mutant allele. Cassette changes by allele conversion can be found on: <http://www.mousephenotype.org/about-ikmc/targeting-strategies>. For example, tm1b allele contains still lacZ reporter cassette, Neo selection cassette is deleted (promotor-driven only).

Please note that these assays are with universal cassette primers other than gene-specific. The confirmation on gene identity performed by e.g. sr genespecific PCR as provided is suggested .

### PCR primer pairs and expected size bands

Assay	Forward Primer	Reverse Primer	Expected Size Band (bp)
lacZ	LacZ_multi_Deen_2F	LacZ_multi_Deen_2R	mut 81 bp,wt without band
Neo	Neo_long_Deen_F1	Neo_long_Deen_R1	mut 186 bp,wt without band

### Primer sequences

Primer Name	Sequence 5' --> 3'
LacZ_multi_Deen_2F	TACTGGAGGCTGAAGTTAGAT
LacZ_multi_Deen_2R	GCGTTTCACCCCTGCCATAA
Neo_long_Deen_F1	TTGAACAAGATGGATTGCACGC
Neo_long_Deen_R1	CCTCGTCCTGCAGTTCAATT

### PCR setup (Qiagen, Hot Start Plus)

### Amplification conditions

Component	Volume (µl)	Final conc.	PCR Settings	Temperature (°C)	Time	# of cycles
DNA (~ 50-100 ng)	2		Denaturation (Melting)	95°C	5 min	1
Q-Solution (5x)	2,5	0,5	Amplification (Melting, An-nealing, Polym.)	94°C	30 sec	
PCR-Buffer (10x)	2,5	1		58°C	45 sec	39
DNTP mix (10 mM)	0,5	0,2		72°C	45 sec	
MgCl <sub>2</sub> (25mM)	1,5	1,5	Polymerisation	72°C	10 min	1
Primer 1 (10 pmol/µl)	1	0,4				
Primer 2 (10 pmol/µl)	1	0,4	Cooling	12°C	hold	1
Taq Polymerase (5 U/µl)	0,3	0,06				
H <sub>2</sub> O	13,7					
Final volume	25					

These PCR conditions have been optimized for our methods and preparation kits. Adoptions may be required.

## Tm1b Allele Conversion PCR-assays

### Allele conversion guide - genotyping tm1b, tm1c and tm1d mice

<http://www.mousephenotype.org/about-ikmc/targeting-strategies>

Tm1b allele is reporter-tagged deletion allele (post-Cre). Critical exon is deleted by creating a frame-shift using Cre method. Neo selection cassette is removed together in promoter-driven strains only. LacZ reporter cassette is kept for visualising gene expression.

Assay	Forward Primer	Reverse Primer	Size Band (bp)	Allele
Tm1b Promotor-driven	tm1b_forw	Floxed LR	380 bp others	tm1b, Promotor-driven tm1a or partially conversion
Flox Promotorless	Floxed PNF	Floxed LR	128 bp ~ 1 kb	tm1b, Promotorless tm1a

### Primer sequences

Primer Name	Sequence 5' --> 3'
tm1b_forw	CGGTCGCTACCATTACCACT
Floxed LR	ACTGATGGCGAGCTCAGACC
Floxed PNF	ATCCGGGGGTACCGCGTCGAG

### PCR setup (Phire Hot Start II)

### Amplification conditions

Component	Volume (µl) 1x	PCR Settings	Temperature (°C)	Time
DNA (~ 50-100 ng)	2,0		1 98°C	30 sec
H <sub>2</sub> O	12,7		2 98°C	5 sec
PCR-Buffer (5x)	4,0		3 58°C	10 sec
DNTP mix (10 mM)	0,4		4 72°C	10 sec
Primer mixed (10 µM)	0,5		5 to 2 + 34 cycles	
Phire Tag (1 U/µl)	0,4		6 72°C	1 min
Final volume	20		7 12°C	hold

These PCR conditions have been optimized for our methods and preparation kits. Adaptations may be required.