

Gene: Otof


Colony prefix: DAQR

Allele: Otof em1(IMPC)Wtsi CRISPR F1 Colony (PM) P1646S

Allele type: Crispr/Cas9 mediated Point mutation

Allele information:

Further information about the allele can be found on the 'International Mouse Phenotyping Consortium' (IMPC) web site at http://www.mousephenotype.org/



Mouse QC information

SNP qPCR	pass	Mutation Sequence confirmed	pass
Mutant Specific SR- PCR	na	Off-target analysis complete	na

Guide RNAs and mutant oligos used in initial experiment

Sequence	Chr	Chr Start	Chr End
CACGGGGCCTTCAGAAATAG	5	30374854	30374873
GACGGCCCCACTTTGGTCCCCATGGGAGAGTGAGGGTTGCC			
AACCGTGTCTTCACGGGGTCTTCAGAAATAGAGGATGAGAAT	5	30374807	30374926
GGTAAGGGACATTGGAAGCAGCTGACTGGCCCTCCT			

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").



Genotyping by end-point PCR

PCRs primer pairs and expected size bands

Assay Type	Assay	Forward Primer	Reverse Primer	Expected Size Band (bp)
Standard PCR	Screening*	Otof_1646-PM_F3	Otof_1646-PM_R3	200

^{*}The screening PCR flanks the SNP region and can be used for sequence verification of the allele. The PCR will not distinguish wild type from mutant mice, however, as a product will be amplified in all cases.

We recommend that mice are sequence-verified with the screening primers to confirm the genotyping qPCR results when establishing the colony, in case of any cross-talk between the assays.

Primer sequences

Primer Name	Primer Sequence (5' > 3')	
Otof_1646-PM_F3	TCCTGACACGCCTCTGTAAA	
Otof_1646-PM_R3	tccccatagagaagtgtgg	

Reaction setup

Reagent	μl
DNA (~50-100 ng)	1
10x Buffer	2
MgCl2 (50 mM)	0.6
Platinum Taq (Invitrogen)	0.2
dNTPs (100 mM)	0.2
Primer 1 (10 μM)	0.4
Primer 2 (10 μM)	0.4
ddH20	15.2
Total	20

Amplification conditions

Step	Conditions	Time
1	94°C	5 min
2	94°C	30 sec
3	58°C	30 sec
4	72°C	1:30 sec
5	Go to '2' + 34 cycles	-
6	72°C	5 min
7	12°C	Forever

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").



Genotyping by SNP qPCR

Primers for LoA qPCR assay

Gene	Source	Forward Primer Seq.	Reverse Primer Seq.	Probe Primer Seq.
Otof	Life	GGTTGCCAACCGTGTCTT	CTGCTTCCAATGTCCCTTACCAT	[VIC]CTCTATTTCTGAAGGCCCCGT
	Technologies	С		[FAM]CTCTATTTCTGAAGACCCCGT

Reactions are performed in a 10µl volume using an Applied Biosystems 7900HT Fast Real-Time PCR System or Applied Biosystems Viia7 with DNA prepared using the Sample-to-SNP[™] kit (Applied Biosystems) from mouse ear biopsies. GTXpress[™] buffer is also used (Applied Biosystems).

Reagent	μl
2x GTXpressTM buffer	5
40x target assay	0.25
ddH2O	3.75
DNA	1

Amplification conditions

Step	Conditions	Time
Pre-read	60°C	30 sec
1	95°C	20 sec
2	95°C	10 sec
3	60°C	30 sec
4	Go to '2' + 34	-
Post-read	60°C	30 sec

Links to information and frequently asked questions

MGP mouse phenotype data:

http://www.mousephenotype.org

How the "critical" exon is decided:

http://www.i-dcc.org/kb/entry/102/

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").



Relevant publications

White, J.K., Gerdin, A.-K., Karp, N.A., Ryder, E., Buljan, M., Bussell, J.N., Salisbury, J., Clare, S., Ingham, N.J., Podrini, C., et al. (2013). Genome-wide Generation and Systematic Phenotyping of Knockout Mice Reveals New Roles for Many Genes. Cell 154, 452–464.

Mali P, Yang L, Esvelt KM, et al (2013) RNA-guided human genome engineering via Cas9. Science 339:823–6. doi: 10.1126/science.1232033

Jinek M, Chylinski K, Fonfara I, et al (2012) A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. Science 337:816–21. doi: 10.1126/science.1225829

Cong L, Ran FA, Cox D, et al (2013) Multiplex genome engineering using CRISPR/Cas systems. Science 339:819–23. doi: 10.1126/science.1231143

Singh P, Schimenti JC, Bolcun-Filas E (2014) A Mouse Geneticist's Practical Guide to CRISPR Applications. Genetics genetics.114.169771—. doi: 10.1534/genetics.114.169771

Brandl C, Ortiz O, Röttig B, et al (2015) Creation of targeted genomic deletions using TALEN or CRISPR/Cas nuclease pairs in one-cell mouse embryos. FEBS Open Bio 5:26–35. doi: 10.1016/j.fob.2014.11.009

Zhou J, Wang J, Shen B, et al (2014) Dual sgRNAs facilitate CRISPR/Cas9 mediated mouse genome targeting. FEBS J. doi: 10.1111/febs.12735

Kraft K, Geuer S, Will AJ, et al (2015) Deletions, Inversions, Duplications: Engineering of Structural Variants using CRISPR/Cas in Mice. Cell Rep. doi: 10.1016/j.celrep.2015.01.016

Shen B, Zhang J, Wu H, et al (2013) Generation of gene-modified mice via Cas9/RNA-mediated gene targeting. Cell Res 23:720–3. doi: 10.1038/cr.2013.46

Wang H, Yang H, Shivalila CS, et al (2013) One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. Cell 153:910–8. doi: 10.1016/j.cell.2013.04.025

Yang H, Wang H, Shivalila CS, et al (2013) One-Step Generation of Mice Carrying Reporter and Conditional Alleles by CRISPR/Cas-Mediated Genome Engineering. Cell 154:1370–1379. doi: 10.1016/j.cell.2013.08.022

This technical data sheet and information ("Datasheet") is supplied by Genome Research Limited ("GRL").